

# Developmental Psychobiology

## **Prior Night Sleep Moderates the Daily Spillover Between Conflict with Peers and Family and Diurnal Cortisol**

Journal:	<i>Developmental Psychobiology</i>
Manuscript ID	DEV-21-090.R3
Wiley - Manuscript type:	Brief Report
Date Submitted by the Author:	n/a
Complete List of Authors:	Armstrong-Carter, Emma; Stanford University, Graduate School of Education Nelson, Benjamin; UNC-Chapel Hill, Department of Psychology & Neuroscience Telzer, Eva; University of North Carolina at Chapel Hill
Keywords:	diurnal cortisol, peer conflict, family conflict, adolescence, daily diary

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17 **Prior Night Sleep Moderates the Daily Spillover**  
18  
19 **Between Conflict with Peers and Family and Diurnal Cortisol**  
20  
21  
22  
23  
24

25  
26 Emma Louise Armstrong-Carter

27  
28 Graduate School of Education, Stanford University  
29

30  
31  
32  
33 Benjamin W. Nelson, Ph.D and Eva H. Telzer, Ph.D

34  
35 Department of Psychology & Neuroscience, University of North Carolina at Chapel Hill  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 Correspondence regarding this manuscript may be addressed to Emma Armstrong-Carter,  
46 [emmaac@stanford.edu](mailto:emmaac@stanford.edu); Stanford University Graduate School of Education, 520 Galvez Mall,  
47 Stanford, CA 94305. The authors have no conflict of interest to declare. Data and syntax are  
48 available upon request.  
49

50  
51 This manuscript was prepared with support from (1) the Institute of Education Sciences  
52 (R305B140009) to Stanford University, awarded to EAC, and the (2) National Institutes of  
53 Health Grant R01DA039923 and National Science Foundation Grant SES 1459719 provided to  
54 EHT, the Department of Psychology at the University of Illinois, and the Department of  
55 Psychology and Neuroscience at the University of North Carolina at Chapel Hill.  
56  
57  
58  
59  
60

## Running Header: SOCIAL CONFLICT, SLEEP AND CORTISOL

## Abstract

We investigated whether daily experiences of conflict with family and peers were associated with fluctuations in diurnal cortisol, and whether sleep buffers the associations between conflict and diurnal cortisol. A racially diverse sample of 370 adolescents (ages 11–18; 57.3% female) provided daily diaries and saliva samples for 4 - 5 days. Hierarchical linear models tested how peer and family conflict were associated with diurnal cortisol (i.e., total cortisol output, slope, and awakening response) the next day, and whether these associations were moderated by sleep duration the previous night. When adolescents experienced peer conflict, they showed higher AUCs the next day if they had slept less the night prior to conflict, but relatively lower CARs and flatter cortisol slopes the next day if they had slept more the night prior to conflict. When adolescents experienced family conflict, they also showed higher AUC the next day if they had slept less the night prior to conflict, but higher CAR the next day if they had slept more the night prior to conflict. Family conflict and sleep were not directly or interactively related to cortisol slope.

1  
2  
3  
4  
5  
6  
7

**Prior Night Sleep Moderates the Daily Spillover**  
**Between Conflict with Peers and Family and Diurnal Cortisol**

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

Close interpersonal relationships are protective for physical and mental health across the lifespan (Holt-Lunstad, Smith, Baker, Harris, & Stephenson, 2015), possibly due to their influence on underlying stress response systems (Nelson, Sheeber, Pfeifer, & Allen, 2020). One candidate biological mechanism influenced by relationships and associated with both mental and physical health outcomes is cortisol — the end product of hypothalamic-pituitary adrenal axis (HPAA) activation. Cortisol has been conceptualized both as a marker of long-term physiological health, and as a reflection of experiences of social adaptation and stress. While healthy cortisol regulation displays diurnal variability that is characterized by a natural rise and fall pattern — peaking shortly after waking and declining throughout late morning into evening — this normative diurnal cortisol rhythm can become dysregulated by adverse social experiences, which may increase risk for mental and physical health challenges, both concurrently and later in life (Adam et al., 2017).

35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Adolescence is a critical developmental transition characterized by heightened socioemotional sensitivity, when youth are increasingly engaging in and valuing social interactions with peers (Dahl, Allen, Wilbrecht, & Suleiman, 2018). Adolescence also involves the recalibration of stress-response systems as a result of acute hormonal changes, which set the stage for life long physical health and well-being (Dahl et al., 2018; Prinstein & Giletta, 2020). In particular, youths' experiences of conflict with peers and family have been associated with variability in their cortisol functioning. For instance, youth who experience peer victimization on average exhibit blunted diurnal cortisol levels, including flattened cortisol awakening response (Jiang et al., 2018; Knack, Jensen-Campbell, & Baum, 2011), flattened diurnal slopes (Jiang et

## SOCIAL CONFLICT, SLEEP AND CORTISOL 3

1  
2  
3 al., 2018), and lower levels of raw diurnal cortisol measured at six time points throughout the  
4  
5 day (Vaillancourt et al., 2008). Youth who experience greater family conflict have shown both  
6  
7 heightened cortisol reactivity at age 7 to 10 (Doom et al., 2018), and blunted cortisol reactivity at  
8  
9 age 14 (Gonzales et al., 2018). Examining whether adolescents' experiences of conflict on one  
10  
11 day predict their cortisol outcomes the next day may offer a more robust predictive model and  
12  
13 reveal previously undetected associations—or spillover—between conflict and cortisol across  
14  
15 days.  
16  
17  
18

19 In addition, sleep promotes adolescents' physical and mental health, and has been directly  
20  
21 linked to cortisol functioning (Heissel, Levy, & Adam, 2017). For instance, both within-subject  
22  
23 and between-subject analyses have shown that longer sleep duration and higher sleep quality is  
24  
25 linked to a steeper decline in cortisol across the day (Ly, McGrath, & Gouin, 2015; Van Lenten  
26  
27 & Doane, 2016; Zeiders, Doane, & Adam, 2011). Moreover, sufficient sleep strengthens youths'  
28  
29 ability to self-regulate their emotions and behaviors, and cope effectively with environmental  
30  
31 stressors (Heissel et al., 2017). For example, poorer sleep functioning exacerbated the positive  
32  
33 association between interparental conflict and young adolescents' aggression (Lemola, Schwarz,  
34  
35 & Siffert, 2012). One study found that family demands were related to a smaller CAR only  
36  
37 among adolescents with poorer sleep (Chiang et al., 2016). This research suggests that adequate  
38  
39 sleep may similarly mitigate the effects of interpersonal conflict on diurnal cortisol functioning,  
40  
41 although this has not yet been explicitly tested. In particular, adequate sleep may enable  
42  
43 adolescents to effectively engage regulatory systems to cope with distress caused by  
44  
45 interpersonal conflict, and thereby buffer any potential negative effects on HPA dysregulation.  
46  
47  
48  
49  
50

51 This study capitalized on the daily diary method to examine (1) How adolescents' daily  
52  
53 experiences of conflict with peers and family relate to their diurnal cortisol functioning the next  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 day; and (2) whether acquiring more sleep the night prior to experiencing conflict mitigates the  
4  
5 daily association between conflict and diurnal cortisol functioning the next day. We examined  
6  
7 three indices of cortisol: *Cortisol Awakening Response* (CAR; the rise in cortisol shortly after  
8  
9 awakening, for which moderate levels are optimal), *Diurnal Cortisol Slope* (the decline in  
10  
11 cortisol across the day, which is optimal when low and steep), and *Area Under the Curve* (AUC;  
12  
13 total cortisol output throughout the day, which is optimal when moderate or low; Chida &  
14  
15 Steptoe, 2009). We hypothesized that conflict with peers and family would be associated with  
16  
17 higher AUC and flatter cortisol slopes the next day when adolescents had slept less than usual,  
18  
19 and not when adolescents slept more than usual, because adequate sleep may facilitate effective  
20  
21 regulation to cope with and reduce physiological stress. We did not hypothesize a direction of  
22  
23 potential associations with CAR, since moderate levels of CAR are linked to optimal physical  
24  
25 and mental health (Stalder et al., 2016).  
26  
27  
28  
29

## 30 31 **Methods**

### 32 33 **Sample and Procedure**

34  
35 Participants were 370 adolescents (57.3% female;  $M_{\text{age}}=14.63$  years,  $SD=1.39$  years;  
36  
37 Range 11–18). The community sample was racially diverse: 39.46% were Non-Latinx White  
38  
39 (N=146), 25.4% Asian (N=94), 17.8% Latinx (N=66), 10.8% African American (N=40), and  
40  
41 6.5% other race (N=24). Approximately 24% of mothers completed high school (23% did not),  
42  
43 and 50% completed postsecondary education (3% declined to answer). Family income ranged  
44  
45 from < \$14,999 to >\$90,000 (Median = \$60,000-\$74,999). Participants were recruited as part of  
46  
47 seven sub-studies in the U.S. Midwest and West between 2012 to 2016. These sub-studies were  
48  
49 all combined for the current sample. All participants were recruited from their community using  
50  
51 convenience sampling, including posting flyers at schools, posting on listservs serving ethnic  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 5

1  
2  
3 minority families, recruiting participants from previous studies who agreed to be contacted for  
4 other research studies, and word of mouth. Participants were compensated \$20 plus a \$20 bonus  
5 for on-time, complete data. Participants were provided with up to 14 days of diary checklists,  
6 which took 5–10 minutes to complete, and a saliva collection kit to complete on days 2-5. The  
7 analyses in the current study only include up to 5 days: days 2-5 when cortisol was collected plus  
8 one day prior (for controlling for prior day levels of the outcome) or after (for testing CAR the  
9 next day). Most participants completed all their dairies (94.90%) and saliva samples (98.92%).  
10 Participants completed their dairies just before bedtime on paper (63.20%) or online (36.80%).  
11 Completion times were verified with an electronic time stamper for paper dairies (Dymo  
12 Corporation, Stamford, CT) and via the website for online dairies. The sponsoring institution's  
13 ethics board approved all study procedures.

### Measures

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1 displays a visual depiction of the timing of measurements. Sleep duration was measured the night prior to conflict with peers and family, which in turn was measured the day before cortisol indices.

**Daily Conflict with Peers and Family.** Participants completed daily dairies to indicate their experiences of family and peer conflict. Each item was coded as 0 = no or 1 = yes. *Family conflict* was the mean of four items: “argued with a sibling”, “got into trouble or were punished by your parents”, “argued with a parent”, and “lied to parent” ( $\alpha = .42$ ; ICC = .35). Relatively low reliability for family conflict may be because each item reflects a unique dimension of family conflict. For instance, adolescents may argue with a sibling, but not with a parent on a given day. *Peer conflict* was the mean of eleven items: “hit, kicked, or shoved a peer”, “threatened, insulted, or made fun of a peer”, “said something mean behind a friend’s back”, “excluded or left a friend

## SOCIAL CONFLICT, SLEEP AND CORTISOL 6

1  
2  
3 out”, “lied to a friend”, “someone online or in a text message threatened, insulted or made fun  
4 of you”, “argued with a friend”, “argued with a boyfriend/girlfriend”, “were excluded or left  
5  
6 out by friends”, “a peer said something mean behind your back”, and “a peer threatened, insulted,  
7  
8 or made fun of you” ( $\alpha = .68$ ;  $ICC = .42$ ).  
9

10  
11  
12 **Sleep Duration.** Adolescents reported on the daily diaries the time they fell asleep the  
13 night before and the time they woke up that morning, yielding a continuous measure of *sleep*  
14 *duration* (in hours). This method is commonly used in diary studies, and is moderately correlated  
15 with sleep estimates from wrist actigraphy (Matthews, Hall, & Dahl, 2014). We time-lagged this  
16 variable so that *sleep duration* reflects sleep the night prior to experiencing conflict ( $M = 8.39$ ,  
17  $SD = 1.74$ , *Range* 0 – 12.5,  $ICC = .18$ ).  
18  
19  
20  
21  
22  
23  
24  
25

26 **Diurnal Cortisol.** Participants provided saliva at four time-points on days 2-5, for a total of  
27 16 samples: (a) immediately upon waking up, (b) 30 min after waking up, (c) 5p.m. (or before  
28 dinner), and (d) 8p.m. (or before bed). Participants were instructed to take their samples before or  
29 > 30 min after brushing teeth, drinking, eating, or using tobacco. Raw cortisol values >60 nmol/L  
30 were excluded as outliers. Participants time-stamped each sample and refrigerated it. The samples  
31 were transferred to the research laboratory and stored in a -80°C freezer, then assayed using high-  
32 sensitivity chemiluminescence- immunoassays (IBL International, Hamburg, Germany). The inter-  
33 assay coefficient of variation was < 8%. Two raw cortisol values exceeding 60 nmol/L were  
34 flagged as outliers and excluded from analyses.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46  
47 *Area under the curve (AUC)* reflects the total daily cortisol output and is positively  
48 associated with chronic stress (Pruessner, 2003). We computed AUC using the trapezoid method  
49 from the first, third, and fourth cortisol measures ( $ICC = .39$ ; Pruessner et al., 2003).  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## SOCIAL CONFLICT, SLEEP AND CORTISOL 7

1  
2  
3 *Cortisol Awakening Response (CAR)* is the steep increase in cortisol from wake to 30–45  
4 min after awakening which mobilizes bodily energy for the demands of the upcoming day (Fries,  
5 Dettenborn, & Kirschbaum, 2009). We computed CAR as the increase in cortisol from wake to  
6 30 min post-wake (ICC = .20; Pruessner et al., 2003); 15–45 minutes between the first two  
7 samples was considered on time. Further information is available in BLINDED, 2021.  
8  
9

10  
11  
12  
13  
14  
15 *Diurnal cortisol slope* represents the decrease in secreted cortisol from morning to  
16 evening. We computed diurnal slopes as the difference between the fourth cortisol sample and  
17 the first morning sample, divided by the time elapsed between these two samples (ICC = .22). A  
18 relatively healthy diurnal slope is typically a steep, negative decline, whereas a flatter (i.e., less  
19 negative) slope is associated with greater stress and cardiovascular risk (Adam et al., 2017). We  
20 time-lagged all cortisol variables so that each reflects cortisol levels the day after adolescents'  
21 experiences of peer and family conflict.  
22  
23  
24  
25  
26  
27  
28  
29

### 30 **Statistical Analyses**

31  
32  
33 Linear mixed effect models nested days (Level 1) within participants (Level 2). We person-  
34 centered all Level 1 predictors, and included intercept person-average values for each of our daily  
35 predictors (Curran & Bauer, 2011). This approach helps to isolate within- vs between- subject  
36 effects. Specifically, daily-spillover can be interpreted more confidently, because the models  
37 control for between-subject differences (Curran & Bauer, 2011). To increase the robustness of our  
38 findings, we additionally controlled for whether the adolescent attended school that day (1 = Yes,  
39 0 = No).  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Model 1 tested family conflict, peer conflict, and sleep duration as simultaneous Level 1  
50 predictors of AUC, CAR and diurnal slope the next day. Model 2 additionally included two daily-  
51 level interaction terms between each daily conflict variable (i.e., family conflict and peer conflict)  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 8

1  
2  
3 and sleep duration the night prior to experiencing conflict. To probe significant interactions, we  
4 used the simple slopes technique at 1SD above and below the mean value of the moderator (Aiken,  
5 West, & Reno, 1991). Missing data were low for family conflict and peer conflict (2.28%) and  
6 sleep duration (4.84%). Missing data were also relatively low for CAR (7.33%), AUC (16.72%)  
7 and slope (13.13%). To manage missing data, we used full information maximum likelihood  
8 (FIML) via the “gsem” command in Stata. We used the observed information matrix (OIM) to  
9 calculate standard errors, which is the default standard error method for the “gsem” command in  
10 Stata (StataSE, Version 15.1.632).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

### 21 Results

22  
23  
24 Table 1 displays descriptive statistics and bivariate correlations. On average, adolescents  
25 slept 8.34 hours per night ( $SD = 1.26$ ). Adolescents experienced at least one type of peer conflict  
26 on 23% of days. Adolescents experienced at least one type of family conflict on 37% of days.  
27  
28  
29

30  
31 Table 2 displays standardized multilevel model regression results. Model 1 demonstrates  
32 direct associations. Family conflict and sleep were not directly related to next day cortisol on  
33 daily or average levels. On the daily level (i.e., within subjects), greater peer conflict was  
34 associated with a relatively flatter cortisol slope the next day, however this association was  
35 qualified by a significant interaction in Model 2, so we interpret it no further.  
36  
37  
38  
39  
40  
41

42 Model 2 reveals five significant daily-level associations between daily conflict and sleep  
43 the prior night predicting next day cortisol indices. First, sleep significantly interacted with peer  
44 conflict and with family conflict to predict AUC the next day. These two interactions display a  
45 consistent pattern. Specifically, as shown in Figure 2 (Panels A and B), peer and family conflict  
46 were only associated with higher AUC the next day when adolescents slept less than their usual  
47 the prior night, but not when adolescents slept more than their usual the night prior to conflict.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 9

1  
2  
3 The simple slope for family conflict was statistically significant ( $p = .024$ ). Although the simple  
4 slope for peer conflict was marginally significant ( $p = .055$ ), it illustrated the same pattern as  
5 family conflict.  
6  
7

8  
9  
10 Second, sleep significantly interacted with peer conflict and with family conflict to  
11 predict CAR the next day. These two interactions displayed opposite patterns for peer conflict  
12 compared to family conflict. Specifically, as shown in Figure 3 (Panel A), peer conflict was  
13 associated with *lower* CAR the next day when adolescents slept less than their usual the prior  
14 night, but not when adolescents slept more than their usual the prior night. However, as shown in  
15 Figure 2 (Panel B), family conflict was associated with *higher* CAR the next day when  
16 adolescents slept less than their usual the prior night, but not when adolescents slept more than  
17 their usual the prior night.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 Third, sleep significantly interacted with peer conflict to predict cortisol slope the next  
29 day. Specifically, as shown in Figure 4, peer conflict was associated with a relatively flatter  
30 cortisol slope the next day when adolescents slept more than usual, but not when they slept less  
31 than usual. Family conflict and sleep were not interactively associated with cortisol slope the  
32 next day. There were no other significant associations in any of these models.  
33  
34  
35  
36  
37  
38  
39

### 40 Discussion

41  
42 Understanding adolescents' diurnal cortisol functioning in the context of their  
43 interpersonal experiences can shed light on how social stressors “get under the skin” during this  
44 critical developmental transition. The physiological embedding of social conflict in turn impacts  
45 physical and mental health throughout the life course. Our study revealed that adolescents  
46 exhibited higher levels of total cortisol output (AUC) the day after experiencing conflict with  
47 peers and family — but only when they had slept less than usual the night prior to conflict. This  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 10

1  
2  
3 consistent pattern suggests that adequate sleep the night before experiencing interpersonal  
4  
5 conflict may mitigate the biological embedding of elevated AUC, which has been associated  
6  
7 with poorer mental and physical health in prior research (Miller, Chen, & Zhou, 2007). Further,  
8  
9 we found that the interplay between adolescents' experiences of conflict and sleep predicting  
10  
11 cortisol awakening response (CAR) and slope occurred only under conditions of adequate sleep,  
12  
13 and depended on the type of conflict. Specifically, adolescents showed lower CAR and flatter  
14  
15 cortisol slope the day after experiencing peer conflict and sleeping more than usual, whereas they  
16  
17 showed higher CAR and no differences in cortisol slope the day after family conflict and  
18  
19 sleeping more than usual. These findings reveal complex daily spillover between interpersonal  
20  
21 conflict, sleep, and multiple indices of HPA function during adolescence.  
22  
23  
24  
25

26 Our primary finding was that adolescents exhibited elevated AUC (i.e., total cortisol  
27  
28 output) the day after they experienced more conflict with peers and family, but only if they slept  
29  
30 less than usual the night before the conflict occurred. In contrast, conflicts with peers and family  
31  
32 were not associated with total cortisol output the next day when adolescents had slept more than  
33  
34 usual. This was a consistent pattern of results for conflict with peers and family. Elevated total  
35  
36 cortisol output is a known risk factor for mental and physical health conditions concurrently and  
37  
38 across the lifespan (Miller, Chen, & Zhou, 2007). When adolescents have slept less than usual,  
39  
40 experiencing interpersonal conflict may be associated with elevated cortisol because  
41  
42 interpersonal conflict is experienced as more stressful, or more physiologically taxing. In  
43  
44 particular, adequate sleep builds adolescents' emotional reserves and may enable them to cope  
45  
46 more effectively with stress they may feel after experiencing conflict with peers and family.  
47  
48  
49  
50  
51 When adolescents sleep enough, they may be able to self-regulate difficult emotions after the  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 11

1  
2  
3 conflict and thereby mitigate potential physiological stress (reflected in elevated total cortisol  
4 output), which would otherwise occur due to the emotional stress.  
5  
6

7  
8 In contrast to AUC, experiences of conflict with peers and family were associated with  
9  
10 CAR only when adolescents slept *more* the previous night. Because inadequate sleep is a strong  
11 predictor of dysregulated CAR (Stalder et al., 2016), it is feasible that interpersonal conflict only  
12 emerges as a predictor of CAR when adolescents have slept enough. Interestingly, and unlike  
13  
14 AUC, the association between conflict and CAR under conditions of adequate sleep diverged for  
15  
16 conflict with peers compared to conflict with family. Specifically, when adolescents had slept  
17  
18 more than usual, they showed *lower* CAR after experiencing more conflict with peers, whereas  
19  
20 they showed *higher* CAR after experiencing more conflict with family. Since moderate levels of  
21  
22 CAR are associated with optimal physical and mental health, it is not entirely clear whether these  
23  
24 reductions and increases in CAR reflect unhealthy blunting or a healthy lowering to optimal  
25  
26 levels (Chida & Steptoe, 2009). On the one hand, conflict is a taxing and stressful experience, so  
27  
28 the reduced CAR after experiencing peer conflict may reflect a negative blunting, which has  
29  
30 been linked to emotional and social difficulties in prior research (Sladek & Doane, 2015). On the  
31  
32 other hand, the elevated CAR observed after family conflict may reflect increased CAR to  
33  
34 unhealthy levels. One study found that family demands were related to lower CAR among  
35  
36 adolescents with poorer sleep (Chiang et al., 2016), which is not inconsistent with our results, but  
37  
38 did not measure family conflict explicitly. In sum, although the directions of associations were  
39  
40 inconsistent, together our findings offer suggestive evidence that conflict may be associated with  
41  
42 alterations or dysregulation in CAR the next day (both higher and lower), when adolescents have  
43  
44 slept more, perhaps because inadequate sleep predicts dysregulated CAR regardless of  
45  
46 interpersonal conflict. Further, our study found intriguing evidence that family and peer conflict  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 may be divergently related to CAR. Future research should clarify the precise mechanisms that  
4  
5 account for these diverging patterns. For example, future work should examine whether the  
6  
7 reductions and elevations in CAR reflect more optimal or less optimal cortisol functioning by  
8  
9 using longitudinal designs to measure the long-term physical and mental implications later in  
10  
11 adolescence and in adulthood.  
12  
13

14  
15 Finally, adolescents also exhibited relatively flatter cortisol slopes the day after they  
16  
17 experienced more conflict with peers, but again only if they slept more than usual the prior night.  
18  
19 Lower, steeper slopes have been associated with more optimal mental and physical health  
20  
21 concurrently and across the lifespan (Adam et al., 2017). When adolescents have slept more than  
22  
23 usual the night before, experiencing conflict with peers may be linked to normative, healthy,  
24  
25 decline in cortisol throughout the day (rather than flattened cortisol slopes), because peer conflict  
26  
27 is experienced as relatively less stressful, or less physiologically taxing, than it would have been  
28  
29 without adequate sleep. When adolescents sleep enough, they may be able to self-regulate  
30  
31 difficult emotions after the conflict and thereby mitigate potential physiological stress (reflected  
32  
33 in flatter cortisol slopes than usual), which would otherwise occur due to the emotional stress. In  
34  
35 this way, experiencing peer conflict after inadequate sleep may be related to a pattern of flat,  
36  
37 sustained cortisol output throughout the day, whereas adequate sleep enables adolescents to cope  
38  
39 more effectively and exhibit lower, more healthy declining cortisol slope throughout the day. In  
40  
41 contrast to peer conflict, family conflict and sleep were not directly or interactively related to  
42  
43 cortisol slope. This is consistent with one prior study that found no association between family  
44  
45 conflict and diurnal cortisol slopes among children age 7 to 9 (Doom et al., 2018). Future  
46  
47 research should further investigate whether peer conflict may have greater salience on  
48  
49 influencing adolescent's cortisol slope, compared to family conflict.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 13

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

There are several directions for future research. First, while it was beyond the scope of our study, future work should examine developmental differences within the adolescent period and early adulthood. Second, our study used subjective reports of sleep duration, which may be influenced by retrospective bias. Future research should use more objective sleep markers such as actigraphy. Third, we were unfortunately unable to control for smoking, alcohol or medication use which could impact cortisol (Adam et al., 2017). Fourth, we calculated CAR from two cortisol samples taken at waking and 30 min after waking; contrasting the more recent recommendation to use two samples taken at 30 and 45 min after waking (Stalder et al., 2016). Fifth, days that are missing from the data (i.e., when adolescents did not respond to the diaries or did not provide cortisol samples) might represent the most difficult days and influence our results.

In sum, our findings suggest that acquiring adequate sleep may mitigate risk for heightened total cortisol output after experiencing social stressors with both peers and family. However, the interplay between conflict and sleep predicting CAR and slope may be more complex and depend on whether the conflict is with peers or family. Future research should investigate the emotional mediators linking peer conflict to cortisol outcomes, as well as long-term associations with physical and emotional health later in life.

## References:

- Adam, E. K., Quinn, M. E., Tavernier, R., McQuillan, M. T., Dahlke, K. A., & Gilbert, K. E. (2017). Diurnal cortisol slopes and mental and physical health outcomes: A systematic review and meta-analysis. *Psychoneuroendocrinology*, *83*(1), 25–41. doi: 10.1016/j.psyneuen.2017.05.018
- Aiken, L., West, S., & Reno, R. (1991). *Multiple regression: Testing and interpreting interactions*. Sage.
- Chiang, J. J., Tsai, K. M., Park, H., Bower, J. E., Almeida, D. M., Dahl, R. E., ... Fuligni, A. J. (2016). Daily family stress and HPA axis functioning during adolescence: The moderating role of sleep. *Psychoneuroendocrinology*, *71*, 43–53. doi: 10.1016/j.psyneuen.2016.05.009
- Chida, Y., & Steptoe, A. (2009). Cortisol awakening response and psychosocial factors: A systematic review and meta-analysis. *Biological Psychology*, *80*(3), 265–278. doi: 10.1016/j.biopsycho.2008.10.004
- Curran, P. J., & Bauer, D. J. (2011). The disaggregation of within-person and between-person effects in longitudinal models of change. *Annual Review of Psychology*, *62*(1), 583–619. doi: 10.1146/annurev.psych.093008.100356
- Dahl, R. E., Allen, N. B., Wilbrecht, L., & Suleiman, A. B. (2018). Importance of investing in adolescence from a developmental science perspective. *Nature*, *554*(7693), 441–450. doi: 10.1038/nature25770
- Doom, J. R., Cook, S. H., Sturza, J., Kaciroti, N., Gearhardt, A. N., Vazquez, D. M., ... Miller, A. L. (2018). Family conflict, chaos, and negative life events predict cortisol activity in



## SOCIAL CONFLICT, SLEEP AND CORTISOL 15

1  
2  
3 low-income children. *Developmental Psychobiology*, 60(4), 364–379. doi:  
4  
5 10.1002/dev.21602  
6

7  
8 Fries, E., Dettenborn, L., & Kirschbaum, C. (2009). The cortisol awakening response (CAR):  
9  
10 Facts and future directions. *International Journal of Psychophysiology*, 72(1), 67–73. doi:  
11  
12 10.1016/j.ijpsycho.2008.03.014  
13

14  
15 Gonzales, N. A., Johnson, M., Shirtcliff, E. A., Tein, J.-Y., Eskenazi, B., & Deardorff, J. (2018).  
16  
17 The role of bicultural adaptation, familism, and family conflict in Mexican American  
18  
19 adolescents' cortisol reactivity. *Development and Psychopathology*, 30(5), 1571–1587.  
20  
21 doi: 10.1017/S0954579418001116  
22

23  
24 Heissel, J. A., Levy, D. J., & Adam, E. K. (2017). Stress, sleep, and performance on standardized  
25  
26 tests: Understudied pathways to the achievement gap. *AERA Open*, 3(3), 1–17. doi:  
27  
28 10.1177/2332858417713488  
29

30  
31 Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and  
32  
33 Social Isolation as Risk Factors for Mortality: A Meta-Analytic Review. *Perspectives on*  
34  
35 *Psychological Science*. doi: 10.1177/1745691614568352  
36

37  
38 Jiang, Y., Li, X., Chen, L., Zhou, G., Zhao, J., & Zhao, G. (2018). Peer victimization and diurnal  
39  
40 cortisol rhythm among children affected by parental HIV: Mediating effects of emotional  
41  
42 regulation and gender differences. *Psychoneuroendocrinology*, 97, 174–181. doi:  
43  
44 10.1016/j.psyneuen.2018.07.010  
45

46  
47 Knack, J. M., Jensen-Campbell, L. A., & Baum, A. (2011). Worse than sticks and stones?  
48  
49 Bullying is associated with altered HPA axis functioning and poorer health. *Brain and*  
50  
51 *Cognition*, 77(2), 183–190. doi: 10.1016/j.bandc.2011.06.011  
52  
53

- 1  
2  
3 Lemola, S., Schwarz, B., & Siffert, A. (2012). Interparental conflict and early adolescents'  
4 aggression: Is irregular sleep a vulnerability factor? *Journal of Adolescence*, *35*(1), 97–  
5 105. doi: 10.1016/j.adolescence.2011.06.001  
6  
7  
8  
9  
10 Ly, J., McGrath, J. J., & Gouin, J.-P. (2015). Poor sleep as a pathophysiological pathway  
11 underlying the association between stressful experiences and the diurnal cortisol profile  
12 among children and adolescents. *Psychoneuroendocrinology*, *57*, 51–60. doi:  
13 10.1016/j.psyneuen.2015.03.006  
14  
15  
16  
17  
18  
19 Matthews, K. A., Hall, M., & Dahl, R. E. (2014). Sleep in healthy Black and White adolescents.  
20 *PEDIATRICS*, *133*(5), e1189–e1196. doi: 10.1542/peds.2013-2399  
21  
22  
23  
24 Nelson, B. W., Pollak, O. H., Clayton, M. G., Telzer, E. H., & Prinstein, M. (2021). *An RDoC-*  
25 *based approach to adolescent self-injurious thoughts and behaviors: The interactive role*  
26 *of social affiliation and cardiac arousal* [Preprint]. PsyArXiv. doi:  
27 10.31234/osf.io/rma69  
28  
29  
30  
31  
32  
33 Nelson, B. W., Sheeber, L., Pfeifer, J., & Allen, N. B. (2020). Psychobiological markers of  
34 allostatic load in depressed and nondepressed mothers and their adolescent offspring.  
35 *Journal of Child Psychology and Psychiatry*, jcpp.13264. doi: 10.1111/jcpp.13264  
36  
37  
38  
39  
40 Nelson, E. E., Jarcho, J. M., & Guyer, A. E. (2016). Social re-orientation and brain development:  
41 An expanded and updated view. *Developmental Cognitive Neuroscience*, *17*, 118–127.  
42 doi: 10.1016/j.dcn.2015.12.008  
43  
44  
45  
46  
47 Prinstein, M. J., & Giletta, M. (2020). Future directions in peer relations research. *Journal of*  
48 *Clinical Child & Adolescent Psychology*, *49*(4), 556–572. doi:  
49 10.1080/15374416.2020.1756299  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SOCIAL CONFLICT, SLEEP AND CORTISOL 17

- 1  
2  
3 Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas  
4  
5 for computation of the area under the curve represent measures of total hormone  
6  
7 concentration versus time-dependent change. *Psychoneuroendocrinology*, *28*(7), 916–  
8  
9 931. doi: 10.1016/S0306-4530(02)00108-7  
10  
11  
12 Pruessner, M., Hellhammer, D. H., Pruessner, J. C., & Lupien, S. J. (2003). Self-reported  
13  
14 depressive symptoms and stress levels in healthy young men: Associations with the  
15  
16 cortisol response to awakening. *Psychosomatic Medicine*, *65*(1), 92–99. doi:  
17  
18 10.1097/01.PSY.0000040950.22044.10  
19  
20  
21 Sladek, M. R., & Doane, L. D. (2015). Daily diary reports of social connection, objective sleep,  
22  
23 and the cortisol awakening response during adolescents' first year of college. *Journal of*  
24  
25 *Youth and Adolescence*, *44*(2), 298–316. doi: 10.1007/s10964-014-0244-2  
26  
27  
28 Somerville, L. H. (2013). The teenage brain: Sensitivity to social evaluation. *Current Directions*  
29  
30 *in Psychological Science*, *22*(2), 121–127. doi: 10.1177/0963721413476512  
31  
32  
33 Stalder, T., Kirschbaum, C., Kudielka, B. M., Adam, E. K., Pruessner, J. C., Wüst, S., ... Clow,  
34  
35 A. (2016). Assessment of the cortisol awakening response: Expert consensus guidelines.  
36  
37 *Psychoneuroendocrinology*, *63*, 414–432. doi: 10.1016/j.psyneuen.2015.10.010  
38  
39  
40 van den Bos, W. (2013). Neural mechanisms of social reorientation across adolescence. *Journal*  
41  
42 *of Neuroscience*, *33*(34), 13581–13582. doi: 10.1523/JNEUROSCI.2667-13.2013  
43  
44  
45 Van Lenten, S. A., & Doane, L. D. (2016). Examining multiple sleep behaviors and diurnal  
46  
47 salivary cortisol and alpha-amylase: Within- and between-person associations.  
48  
49 *Psychoneuroendocrinology*, *68*, 100–110. doi: 10.1016/j.psyneuen.2016.02.017  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Zeiders, K. H., Doane, L. D., & Adam, E. K. (2011). Reciprocal relations between objectively  
4 measured sleep patterns and diurnal cortisol rhythms in late adolescence. *Journal of*  
5  
6  
7 *Adolescent Health, 48*(6), 566–571. doi: 10.1016/j.jadohealth.2010.08.012  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## SOCIAL CONFLICT, SLEEP AND CORTISOL 19

Table 1

Descriptive statistics and bivariate correlations among study constructs based on 370 adolescents.

	1	2	3	4	5	6	7	8	9
1 Daily Family Conflict	1								
2 Daily Peer Conflict	0.32***	1							
3 Sleep Duration	-0.01	-0.14**	1						
4 CAR	0.05	-0.02	-0.13*	1					
5 AUC	0.00	-0.02	-0.01	-0.06	1				
6 Slope	-0.01	0.00	-0.05	0.26***	-0.63***	1			
7 Female	0.03	0.15**	-0.03	0.06	0.09	-0.05	1		
8 Age	-0.12*	0.07	-0.32***	-0.08	-0.31***	0.30***	0.06	1	
9 Maternal Education	-0.07	-0.15**	0.17**	0.022	0.07	-0.02	0.04	-0.19***	1
<i>Mean</i>	0.14	0.05	8.34	4.48	160.64	-1.27	0.57	0.05	3.73
<i>SD</i>	0.16	0.10	1.26	9.54	68.37	0.74	0.50	1.09	1.92
<i>Min</i>	0.00	0.00	2.00	-27.48	-185.58	-4.51	0.00	-2.07	0.00
<i>Max</i>	0.88	1.00	11.75	41.60	492.92	1.18	1.00	2.70	6.00

Table 2

Multi-Level Standardized Regression Results based on a sample of 370 adolescents.

	Model 1						Model 2					
	Next Day CAR		Next Day AUC		Next Day Slope		Next Day CAR		Next Day AUC		Next Day Slope	
	<i>B</i>	<i>SE</i>	<i>B</i>	<i>SE</i>	<i>B</i>	<i>SE</i>	<i>B</i>	<i>SE</i>	<i>B</i>	<i>SE</i>	<i>B</i>	<i>SE</i>
Constant	0.340	(0.365)	0.171	(0.374)	0.661	(0.380)	0.413	(0.362)	0.200	(0.374)	0.708	(0.375)
School Day	0.076	(0.076)	<b>0.269</b>	<b>(0.076)***</b>	<b>-0.191</b>	<b>(0.077)*</b>	0.089	(0.075)	<b>0.282</b>	<b>(0.076)***</b>	<b>-0.182</b>	<b>(0.077)*</b>
Daily Family Conflict	0.225	(0.217)	0.219	(0.214)	0.197	(0.222)	0.203	(0.216)	0.267	(0.213)	0.182	(0.223)
Daily Peer Conflict	-0.818	(0.491)	0.299	(0.515)	<b>-1.761</b>	<b>(0.507)***</b>	-0.721	(0.485)	0.313	(0.511)	<b>-1.675</b>	<b>(0.506)***</b>
Prior Night Sleep	-0.002	(0.026)	-0.002	(0.025)	-0.016	(0.026)	0.003	(0.026)	-0.002	(0.025)	-0.012	(0.026)
Average Family Conflict	0.112	(0.337)	-0.001	(0.358)	0.190	(0.363)	0.101	(0.334)	0.082	(0.358)	0.194	(0.359)
Average Peer Conflict	-0.503	(0.565)	-0.524	(0.594)	-1.425	(0.783)	-0.595	(0.560)	-0.621	(0.595)	-1.511	(0.774)
Average Sleep	-0.041	(0.042)	-0.032	(0.044)	-0.067	(0.044)	-0.051	(0.042)	-0.037	(0.044)	-0.073	(0.044)
Daily Peer Conflict X Daily Sleep							<b>-1.437</b>	<b>(0.406)***</b>	<b>-0.845</b>	<b>(0.406)*</b>	<b>-0.861</b>	<b>(0.417)*</b>
Daily Family Conflict X Daily Sleep							<b>0.545</b>	<b>(0.165)***</b>	<b>-0.330</b>	<b>(0.164)*</b>	0.310	(0.171)

Note: Significant associations are bolded. \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ . Sleep = Sleep Duration. CAR = Cortisol Awakening Response. AUC = Area Under the Curve (i.e., total cortisol output). Slope = Diurnal Cortisol Slope. "Daily" variables are person-mean centered, whereas "average" variables reflect person-average values.

## SOCIAL CONFLICT, SLEEP AND CORTISOL 21

Figure 1

Summary of measurement timing. Sleep duration was measured the night prior to conflict with peers and family, which in turn was measured the day before cortisol indices.





Figure 2

Experiencing conflict with peers and family predicted higher total cortisol output (AUC) the next day only when adolescents slept less than usual the night prior to conflict.





## SOCIAL CONFLICT, SLEEP AND CORTISOL 23

Figure 3

Peer conflict predicted lower CAR the next day, but family conflict predicted higher CAR the next day, when adolescents slept more than usual.




Figure 4

Peer conflict was associated with a flatter than usual cortisol slope the next day when adolescents slept more than usual, but not when they slept less than usual.

